



勝利動物醫院
Victory Animal Hospital

Victory Animal Hospital

Blk B, 1/F, Chung Yuen Mansion, 71b Waterloo Rd., Kln, Hong Kong

Phone: 852 2711 0332

Email: info@globavet.com

Cruciate Ligament Rupture in Dogs

What and where are the cruciate ligaments?

The cruciate ligaments are two bands of fibrous tissue located within each knee joint. They join the femur and tibia (the bones above and below the knee joint) together so that the knee works as a hinged joint.

One ligament runs from the inside to the outside the knee joint and the other from the outside to the inside, crossing over each other in the middle. In the dog, the ligaments are called the cranial and caudal cruciate ligament. In dogs, the most common knee injury is a rupture of the cranial cruciate ligament.


How does a cranial cruciate injury occur?

Based on its anatomy, the knee joint is a "hinge joint". It is relatively unstable because there are no interlocking bones in the joint. Instead it is held together by several ligaments, including the cruciate ligaments, which allow it to move back and forth like a hinge, but restrict its side-to-side motion.

Acute or traumatic cruciate rupture is caused by a twisting injury to the knee joint. This occurs most often when the dog is running and suddenly changes direction. This places the majority of the body weight on the knee joint, and excessive rotational and shearing forces are placed on the cruciate ligaments. This injury usually affects the anterior or cranial (front) ligament. A cruciate ligament rupture is usually extremely painful and the knee joint becomes unstable, resulting in lameness.

Obese dogs appear to be more predisposed to developing a cruciate rupture. In these dogs, the injury may occur with minor trauma to the knee. Dogs with other knee problems such as a luxating patella may also be predisposed to rupturing their cruciate ligaments.

A more chronic form of cruciate damage can occur due to progressive weakening of the ligaments as a result of repeated trauma or arthritic disease. Initially, the ligament becomes stretched or partially torn and lameness may be only slight and intermittent. With continued use of the joint, the condition gradually gets worse until a complete rupture occurs.



Used Under License
Copyright© Lifelearn Inc.

How is it diagnosed?

During the lameness examination, the veterinarian will try to demonstrate a particular movement, called a *cranial or anterior drawer sign*. This abnormal forward movement of the tibia (lower leg bone) in front of the femur (thigh bone) indicates laxity in the knee joint. It may be necessary to administer a sedative to relax the dog enough that the veterinarian can perform this test. Other diagnostic tests such as radiographs (x-rays) may also be necessary.



Used Under License
Copyright © Lifelearn Inc.

Is other joint damage common?


Inside the knee joint are pieces of cartilage called *menisci*. The menisci act as shock absorbers between the femur and tibia. The menisci are often damaged when the cruciate ligaments rupture. They are usually repaired at the same time as the ligament surgery.

Is an operation always necessary?

Unfortunately, most dogs will eventually require surgery to correct this painful injury.

What does surgery involve?

There are various surgical techniques performed to stabilize the knee joint following cruciate rupture. There is a newer surgical technique available called CORA based leveling osteotomy (CBLO) that is especially beneficial for larger, more athletic dogs. Your veterinarian will discuss the best treatment option for your pet.



Used Under License
Copyright © Lifelearn Inc.

Is post-operative care difficult?

It is important that your dog have limited activity for six to eight weeks after surgery. The activity restriction may be less if a CBLO has been performed. Provided you are able to carry out your veterinarian's instructions, good function should return to the limb within three months.

Unfortunately, regardless of the technique used to stabilize the joint, arthritis is likely to develop in the joint as your dog ages. Arthritis develops more slowly and to a lesser degree if the dog undergoes surgery. Weight control and nutritional supplements such as glucosamine / chondroitin may help delay the onset of arthritis in your pet. Many dogs will receive physical therapy after the surgery to speed recovery and reduce complications. Your veterinarian will discuss your pet's recommended post-operative care with you prior to surgery.

Is obesity such a problem?

Obesity or excess weight can predispose dogs to cruciate ligament rupture. If your dog is overweight, the post-surgical recovery time will be much longer. Obesity also greatly increases the risk of injury to the other knee, especially during the recuperation period. Weight loss is as important as surgery in ensuring rapid return to normal function, as well as being a preventive measure to help protect your dog against this debilitating injury.

This client information sheet is based on material written by: Ernest Ward, DVM

© Copyright 2009 Lifelearn Inc. Used and/or modified with permission under license.